arXiv:hep-th/9612236v1 26 Dec 1996

SMI-25-96

Incidence Matrix Description of Intersecting p-brane Solutions

I.Ya.Aref'eva [*]
Steklov Mathematical Institute,
Vavilov 42, GSP-1, 117966, Moscow, Russia
and
O.A.Rytchkov [†]
Department of Theoretical Physics,
Moscow State University,
Moscow 119 899, Russia

Abstract

An algebraic method for a general construction of intersecting p-brane solutions in diverse spacetime dimensions is discussed. An incidence matrix describing configurations of electric and magnetic fields is introduced. Intersecting p-branes are specified by solutions of a system of characteristic algebraic equations for the incidence matrix. This set of characteristic equations generalizes a single characteristic equation found before for a special "flower" ansatz. The characteristic equations admit solutions only for quantized values of scalar coupling parameters. A wide list of examples including solutions with regular horizons and non-zero entropy in D=11 and D=10 theories is presented.

---

[*] e-mail: arefeva@arevol.mian.su

[†] e-mail: rytchkov@grg1.phys.msu.su

1 Introduction

Soliton solutions in $D = 10$ superstring theories and $D = 11$ supergravity support an idea that five known $D = 10$ superstrings are in fact five different points of an unique M-theory [1]-[4]. In this context it is interesting to understand whether an existence of soliton solutions is a privilege of effective superstring theories or $D = 11$ supergravity. For this purpose it is useful to have a list of soliton solutions in D-dimensional theory with the following action

$$I = \frac{1}{2\kappa^2}\int d^Dx\sqrt{-g}(R - \frac{1}{2}(\nabla\phi)^2 - \frac{e^{-\alpha\phi}}{2(d+1)!}F^2_{d+1}), \tag{1.1}$$

where F_{d+1} is a $d+1$ differential form, $F_{d+1} = d\mathcal{A}_d$, ϕ is a dilaton. In general, the fields F and ϕ are linear combinations of a large variety of gauge fields and scalars.

The aim of this paper is to present a systematic description of a class of soliton solutions in higher dimensional gravity coupled with matter. This class describes multi-soliton composite p-brane solutions. Elementary p-brane solutions have been found years ego [5]-[9] (see also [10]-[13]). The recent microscopic interpretation of the Bekenstein-hawking entropy within string theory [15]-[17] has stimulated an investigation of p-brane solutions [18]-[37]. Composite p-brane solutions for $D = 11$ and $D = 10$ have been obtained recently [18, 20, 19]. We will also search for an universal higher dimensional (higher then D=11) theory which has p-brane solutions that after dimension reduction could produce all known $D = 11$ and $D = 10$ p-brane solutions.

To get a wide class of composite p-brane solutions with non-zero electric and magnetic charges in theory (1.1) we will use an algebraic method [29, 33, 37], which permits to reduce the problem of finding solutions to an algebraic problem.

We will describe the class of solutions with a $N + 2$-blocks metric and a gauge field $\mathcal{A}$ which has E electric and M magnetic components (branches)

$$\mathcal{A} = \sum_a^E \mathcal{A}_a^{(\mathcal{E})} + \sum_b^M \mathcal{A}_b^{(\mathcal{M})}. \tag{1.2}$$

An $N + 2$-blocks metric has the following form

$$ds^2 = e^{2A(x)}\eta_{\mu\nu}dy^\mu dy^\nu + \sum_{i=1}^N e^{2F_i(x)}dz_i^{m_i}dz_i^{m_i} + e^{2B(x)}dx^\gamma dx^\gamma, \tag{1.3}$$

where $\eta_{\mu\nu}$ is a flat Minkowski metric, μ, ν run from 0 to $q-1$, m_i runs from 1 to r_i, γ runs from 1 to $s+2$ and

$$D = q + \sum_i^N r_i + s + 2, \tag{1.4}$$

In (1.3) A, B and F_i are functions of x.

To describe electric (magnetic) configurations $\mathcal{A}_a^{(\mathcal{E})}$ ($\mathcal{A}_b^{(\mathcal{M})}$) we will introduce an electric (magnetic) *incidence* matrix. The incidence matrix, say the electric incidence matrix, is an rectangular $N \times E$ matrix,

$$\Delta = (\Delta_{ai}), \quad a = 1, ...E, \quad i = 1, ...N, \tag{1.5}$$

which rows correspond to independent branches of electric gauge field and columns refer to the number of the blocks in the metric (1.3). The entries of the incidence matrix are equal to 0 or 1. Field configurations $\mathcal{A}_a^{(\mathcal{E})}$ will be in the following form

$$\mathcal{A}_a^{(\mathcal{E})} = h_a H_a^{-1} \omega_0 \prod_{\{i|\Delta_{ai}=1\}} \wedge \omega_i, \tag{1.6}$$

where ω_0 and ω_i are q and r_i-form, respectively,

$$\omega_0 = dy^0 \wedge \ldots \wedge dy^{q-1}, \quad \omega_i = dz_i^1 \wedge \ldots \wedge dz_i^{r_i}.$$

H_a are some functions of x-variables, $a = 1, \ldots, E$.

We will show that in the case of a pure electric ansatz,

$$\mathcal{A} = \sum_a^E \mathcal{A}_a^{(\mathcal{E})}, \tag{1.7}$$

the following metric

$$ds^2 = (H_1 H_2 ... H_E)^{-2t\sigma} \sum_{\mu,\nu=0}^{q-1} \eta_{\mu\nu} dy^\mu dy^\nu +$$

$$(H_1 H_2 ... H_E)^{2u\sigma} [\sum_i (\prod_a H_a^{-\Delta_{ai}})^\sigma \sum_{m_i=1}^{r_i} dz_i^{m_i} dz^{m_i} + \sum_\gamma dx^\gamma dx^\gamma], \tag{1.8}$$

is a solution of the theory (1.1). Here

$$u = \frac{d}{2(D-2)}, \quad t = \frac{1}{2} - u, \quad \sigma = \frac{1}{dt + \alpha^2/4}, \tag{1.9}$$

$H_1, H_2 ..., H_E$ are arbitrary harmonic functions of x-variables and the incindence matrix Δ satisfies the following $E(E-1)/2$ *characteristic* equations

$$(q + \frac{\alpha^2}{2}) - \frac{d^2}{D-2} + \sum_{i=1}^N r_i \Delta_{ai} \Delta_{a'i} = 0, \quad a \neq a'. \tag{1.10}$$

Note that the characteristic equations (1.10) generalize a characteristic algebraic equation for a so-called "flower' ansatz [37]. In this ansatz $\sum_{i=1}^N r_i \Delta_{ai} \Delta_{a'i} = 0$, and one has a single equation [37]

$$(q + \frac{\alpha^2}{2}) = \frac{d^2}{D-2}, \tag{1.11}$$

In the case when electric and magnetic components of the d-form gauge field are non-zero (see eqs. (2.2) below) the metric solving the theory (1.1) has the form

$$ds^2 = (H_1^{-1} H_2^{-1} ... H_E^{-1} U_1 U_2 ... U_M)^{2t\sigma} (H_1 H_2 ... H_E)^\sigma \times$$

$$\{(H_1 H_2 ... H_E U_1 U_2 ... U_M)^{-\sigma} \sum_{\mu,\nu=0}^{q-1} \eta_{\mu\nu} dy^\mu dy^\nu +$$

$$(U_1U_2...U_M)^{-\sigma}\sum_i(\prod_a H_a^{-\Delta_{ai}}\prod_b U_b^{\Lambda_{bi}})^{\sigma}\sum_{m_i=1}^{r_i} dz_i^{m_i}dz_i^{m_i}+\sum_\alpha dx^\alpha dx^\alpha\}, \qquad (1.12)$$

where H_a and U_b are arbitrary harmonic functions of x-variables, $a = 1, .., E$, $b = 1, .., M$, and the electric and magnetic incidence matrix, Δ_{ai} and Λ_{bi}, satisfy two more conditions

$$s+\frac{\alpha^2}{2}-\frac{d^2}{D-2}+\sum_{i=1}^{N} r_i\Lambda_{bi}\Lambda_{b'i}=0, b\neq b', \quad b,b'=1,..M \qquad (1.13)$$

$$\frac{\alpha^2}{2}-\frac{d^2}{D-2}+\sum_{i=1}^{N} r_i\Delta_{ai}\Lambda_{bi}=0, \quad a=1,...E, \quad b=1,..M \qquad (1.14)$$

The characteristic equations admit solutions only for quantized values of scalar coupling parameters that is in accordance with [29, 30, 37]. Note that the case when $\alpha = 0$ (D=11 supergravity) p-brane solutions are given by the above formula with $\alpha = 0$. These equations are very restrictive since they must be solved for integers and they admit non-trivial solutions only for special D. The selected D are the same as for the "flower" ansatz [37]. In particular, they are $D = 10,\ 11,\ 18,\ 20,\ 26$.

For the case of two (k) gauge fields we will introduce four ($2k$) incidence matrices and the corresponding expressions for metric will be very similar (see eqs.(3.23) below). The corresponding incidence matrix will satisfy to relations similar to (1.10), (1.13) and (1.14).

The algebraic method [29, 33, 37] can be also used for finding solutions with depending harmonic functions. In particular, one can consider a generalized dyonic ansatz that contain equal numbers of magnetic and electric branches, $\mathcal{E} = \mathcal{M}$ and the same set of harmonic functions $H_i = U_i$, $i = 1, ...\mathcal{E}$. This construction gives in $D = 11$ a new dyonic solution with regular horizon and non-zero entropy.

The paper is organized as follows. In Section 2 we present soliton solutions for the theory (1.1) with one d-form field with dilaton and in Section 3 we present soliton solutions for a theory with two antisymmetric fields and dilaton. Both sections start from general considerations and are concluded by some examples. There one can also find relations with known solutions. In section 4 we collect some examples of solutions of the form (1.12) with depending functions $H_1, H_2...H_E$ and $U_1U_2...U_M$.

2 Gravity Coupled with d-form and Dilaton

2.1 General Consideration

Let M^D be a manifold with a following structure:

$$M^D = \mathbf{M}^q\times\mathbf{R}^{r_1}\times\cdots\times\mathbf{R}^{r_N}\times\mathbf{R}^{s+2},$$

and the metric has the form (1.3).

For the d-form we are going to consider a class of ansatzes corresponding to E electric and M magnetic charges. Each ansatz we will connect with two rectangular matrices $\Delta_{ai},\ a = 1..E,\ i = 1..N$ and $\Lambda_{bi},\ b = 1..M,\ i = 1..N$ with the following properties:

1) $\Delta_{ai} = 1$ or 0;

1') $\Lambda_{bi} = 1$ or 0;

2) there are no a_1 and a_2 ($a_1 \neq a_2$) such that: $\Delta_{a_1 i} = \Delta_{a_2 i}$ for all i

2') there are no b_1 and b_2 ($b_1 \neq b_2$) such that: $\Lambda_{b_1 i} = \Lambda_{b_2 i}$ for all i;

3) there are no a_1 and a_2 ($a_1 \neq a_2$) and $i_1,\ i_2$ such that: $\Delta_{a_1 i} = \Delta_{a_2 i}$ for all $i \neq i_1,\ i_2$ and $r_{i_1} = r_{i_2} = 1$

3') there are no b_1 and b_2 ($b_1 \neq b_2$) and $i_1,\ i_2$ such that: $\Delta_{b_1 i} = \Delta_{b_2 i}$ for all $i \neq i_1,\ i_2$ and $r_{i_1} = r_{i_2} = 1$

4) $\sum_{i=1}^{N} r_i \Delta_{ai} = d - q$ for all a;

4') $\sum_{i=1}^{N} r_i \Lambda_{bi} = d - s$ for all b;

5)$\Delta_{ai} \neq \Lambda_{bi}$ if $s = 1,\ q = 1$

An "electric" field is assumed in the form

$$\mathcal{A} = \sum_{a=1}^{\mathcal{E}} h_a e^{C_a(x)} \omega_0^q \prod_{\{i|\Delta_{ai}=1\}} \wedge\, \omega_i \tag{2.1}$$

where ω_0^q ia a q-form

$$\omega_0^q = dy^0 \wedge \ldots \wedge dy^{q-1}$$

ω_i is a r_i-form

$$\omega_i = dz_i^1 \wedge \ldots \wedge dz_i^{r_i}.$$

A "magnetic" field is assumed in the form

$$F = \sum_{b=1}^{M} v_b e^{\alpha\phi + \chi_b(x)} *_b\, d\chi_b(x), \tag{2.2}$$

where $*_b$ is a Hodge dual on

$$\mathbf{M}_b^{d+2} = (\prod_{\{i|\Lambda_{bi}=1\}} \mathbf{R}^{r_i} \times) \mathbf{R}^{s+2}.$$

We will refer to the different terms in (2.1) and (2.2) as to the different branches of electric and magnetic fields. In order to simplify the Ricci tensor [33, 29, 37] we assume

$$qA + \sum r_i F_i + sB = 0. \tag{2.3}$$

We get

$$R_{\mu\nu} = -\eta_{\mu\nu} e^{2(A-B)} \Delta A, \tag{2.4}$$

$$R_{m_i n_i} = -\delta_{m_i n_i} e^{2(F_i - B)} \Delta F_i, \tag{2.5}$$

$$R_{\alpha\beta} = -q\partial_\alpha A \partial_\beta A - \sum_i r_i \partial_\alpha F_i \partial_\beta F_i - s\partial_\alpha B \partial_\beta B - \delta_{\alpha\beta} \Delta B. \tag{2.6}$$

Components of the energy-momentum tensor for the d-form gauge field,

$$T_{MN}^{(F)} = \frac{1}{2d!}(F_{MM_1\ldots M_d} F_N^{M_1\ldots M_d} - \frac{1}{2(d+1)} g_{MN} F^2) e^{-\alpha\phi} \tag{2.7}$$

for the above ansatz are

$$T_{\mu\nu}^{(F)} = -\eta_{\mu\nu} e^{2(A-B)} [\sum_{a=1}^{E} \frac{h_a^2}{4} e^{-\alpha\phi - 2qA - 2\sum_{i=1}^{N} \Delta_{ai} r_i F_i + 2C_a} (\partial C_a)^2 +$$

$$+\sum_{b=1}^{M}\frac{v_b^2}{4}e^{\alpha\phi+2sB+2\sum_{i=1}^{N}\Lambda_{bi}r_iF_i+2\chi_b}(\partial\chi_b)^2], \tag{2.8}$$

$$T^{(F)}_{m_in_i}=\delta_{m_in_i}e^{2(F_i-B)}[\sum_{a=1}^{E}\frac{h_a^2}{4}e^{-\alpha\phi-2qA-2\sum_{i=1}^{N}\Delta_{ai}r_iF_i+2C_a}(\partial C_a)^2(1-2\Delta_{ai})-$$

$$-\sum_{b=1}^{M}\frac{v_b^2}{4}e^{\alpha\phi+2sB+2\sum_{i=1}^{N}\Lambda_{bi}r_iF_i+2\chi_b}(\partial\chi_b)^2(1-2\Lambda_{bi})], \tag{2.9}$$

$$T^{(F)}_{\alpha\beta}=-\sum_{a=1}^{E}\frac{h_a^2}{2}e^{-\alpha\phi-2qA-2\sum_{i=1}^{N}\Delta_{ai}r_iF_i+2C_a}[\partial_\alpha C_a\partial_\beta C_a-\frac{\delta_{\alpha\beta}}{2}(\partial C_a)^2]-$$

$$\sum_{b=1}^{M}\frac{v_b^2}{2}e^{\alpha\phi+2sB+2\sum_{i=1}^{N}\Lambda_{ai}r_iF_i+2\chi_b}[\partial_\alpha\chi_b\partial_\beta\chi_b-\frac{\delta_{\alpha\beta}}{2}(\partial\chi_b)^2]. \tag{2.10}$$

All others components are zero. Note that we assume 3) and 3') to guarantee the diagonal form of the energy-momentum tensor.

Components of the energy-momentum tensor for dilaton

$$T^{(\phi)}_{MN}=\frac{1}{2}(\partial_M\phi\partial_N\phi-\frac{1}{2}g_{MN}(\nabla\phi)^2)$$

have the following non-zero components

$$T^{(\phi)}_{\mu\nu}=-\frac{1}{4}\eta_{\mu\nu}e^{2(A-B)}(\partial\phi)^2, \tag{2.11}$$

$$T^{(\phi)}_{m_in_i}=-\frac{1}{4}\delta_{m_in_i}e^{2(F_i-B)}(\partial\phi)^2, \tag{2.12}$$

$$T^{(\phi)}_{\alpha\beta}=\frac{1}{2}[\partial_\alpha\phi\partial_\beta\phi-\frac{1}{2}\delta_{\alpha\beta}(\partial\phi)^2]. \tag{2.13}$$

If we assume the following "no-force" conditions

$$-\alpha\phi-2qA-2\sum_{i=1}^{N}\Delta_{ai}r_iF_i+2C_a=0,\ a=1..E \tag{2.14}$$

$$\alpha\phi+2sB+2\sum_{i=1}^{N}\Lambda_{bi}r_iF_i+2\chi_b=0,\ b=1..M \tag{2.15}$$

the form of T_{MN} crutially simplifies and the Einstein equations cast into the form

$$\Delta A=\sum_{a=1}^{E}th_a^2(\partial C_a)^2+\sum_{b=1}^{M}uv_b^2(\partial\chi_b)^2, \tag{2.16}$$

$$\Delta F_i=\sum_{a=1}^{E}h_a^2(\partial C_a)^2(u-\frac{1}{2}\Delta_{ai})-\sum_{b=1}^{M}v_b^2(\partial\chi_b)^2(u-\frac{1}{2}\Lambda_{bi}), \tag{2.17}$$

$$-q\partial_\alpha A\partial_\beta A-\sum_i r_i\partial_\alpha F_i\partial_\beta F_i-s\partial_\alpha B\partial_\beta B-\delta_{\alpha\beta}\Delta B$$

$$=\frac{1}{2}\partial_\alpha\phi\partial_\beta\phi-\sum_{a=1}^{E}h_a^2[\frac{1}{2}\partial_\alpha C_a\partial_\beta C_a-u\delta_{\alpha\beta}(\partial C_a)^2-$$

$$\sum_{b=1}^{M} v_b^2[\frac{1}{2}\partial_\alpha\chi_b\partial_\beta\chi_b - t\delta_{\alpha\beta}(\partial\chi_b)^2], \tag{2.18}$$

where t and u are given by (1.9).

Equation of motion for dilaton reads

$$\triangle\phi - \frac{\alpha}{2}[\sum_{a=1}^{E} h_a^2(\partial C_a)^2 - \sum_{b=1}^{M} v_b^2(\partial\chi_b)^2] = 0 \tag{2.19}$$

From Maxwell's equations under conditions (2.14) and (2.15) we get

$$\Delta C_a = (\partial C_a)^2 \tag{2.20}$$

From Bianchi identity for F under conditions (2.14) and (2.15) we have

$$\Delta\chi_b = (\partial\chi_b)^2, \tag{2.21}$$

therefore, $H_a = e^{-C_a}$, $a = 1,..,E$ and $U_b = e^{-\chi_b}$, $b = 1,...M$ are harmonic functions.

We solve (2.16), (2.17), (2.19) and the diagonal part of (2.18) supposing

$$A = \sum_{a=1}^{E} th_a^2C_a + \sum_{b=1}^{M} uv_b^2\chi_b \tag{2.22}$$

$$F_i = \sum_{a=1}^{E} h_a^2C_a(\frac{1}{2}\Delta_{ai} - u) - \sum_{b=1}^{M} v_b^2\chi_b(\frac{1}{2}\Lambda_{bi} - u) \tag{2.23}$$

$$B = -u\sum_{a=1}^{E} h_a^2C_a - t\sum_{b=1}^{M} v_b^2\chi_b \tag{2.24}$$

$$\phi = \frac{\alpha}{2}[\sum_{a=1}^{E} h_a^2C_a - \sum_{b=1}^{M} v_b^2\chi_b] \tag{2.25}$$

But now we have to guarantee relations (2.3), (2.14), (2.15) and non-diagonal part of (2.18). Let us analyze these conditions step by step.

A substitution of (2.22)–(2.25) in (2.14) under an assumption of the independence of C_a and χ_b gives three types of relations

$$1 = \frac{\alpha^2}{4}h_a^2 + qth_a^2 + \sum_{i=1}^{N} r_i\Delta_{ai}h_a^2(\frac{1}{2}\Delta_{ai} - u) \tag{2.26}$$

$$0 = \frac{\alpha^2}{4} + qt + \sum_{i=1}^{N} r_i\Delta_{ai}h_a^2(\frac{1}{2}\Delta_{a'i} - u),\ a \neq a' \tag{2.27}$$

$$0 = -\frac{\alpha^2}{4} + qu - \sum_{i=1}^{N} r_j\Delta_{ai}(\frac{1}{2}\Lambda_{bi} - u) \tag{2.28}$$

Taking into account requirement 4) on the electric incidence matrix from equation (2.26) we get

$$h_a^2 = h^2 = \sigma \tag{2.29}$$

where σ is given by equation (1.9).

Equations (2.27) and (2.28) together with conditions 4) and 4') give

$$(D-2)(\frac{\alpha^2}{2}+q)-d^2+(D-2)\sum_{i=1}^{N}r_i\Delta_{ai}\Delta_{a'i}=0,\ a\neq a' \tag{2.30}$$

$$(D-2)\frac{\alpha^2}{2}-d^2+(D-2)\sum_{i=1}^{N}r_i\Delta_{ai}\Lambda_{bi}=0, \tag{2.31}$$

that are nothing but conditions (1.10) and (1.14) mentioned in the Introduction.

A substitution of (2.22)–(2.25) in (2.15) under an assumption of the independence of C_a and χ_b gives once again three types of relations. They are

$$1=\frac{\alpha^2}{4}v_b^2+stv_b^2+\sum_{i=1}^{N}r_i\Lambda_{bi}v_b^2(\frac{1}{2}\Lambda_{bi}-u) \tag{2.32}$$

$$0=\frac{\alpha^2}{4}+st+\sum_{i=1}^{N}r_i\Lambda_{bi}(\frac{1}{2}\Lambda_{b'i}-u),\ b\neq b' \tag{2.33}$$

$$0=\frac{\alpha^2}{4}-su+\sum_{i=1}^{N}r_i\Lambda_{bi}(\frac{1}{2}\Delta_{ai}-u) \tag{2.34}$$

From (2.32) it follows that

$$v_b^2=v^2=\sigma \tag{2.35}$$

The condition (2.33) gives

$$(D-2)(\frac{\alpha^2}{2}+s)-d^2+(D-2)\sum_{i=1}^{N}r_i\Lambda_{bi}\Lambda_{b'i}=0, \tag{2.36}$$

that is nothing but (1.13). The relation (2.34) coincides with (2.31).

Substituting (2.22)–(2.25) in (2.3) one can check that equation (2.3) is fulfilled. Indeed, a contribution of terms containing C_a in the LHS of (2.3) is

$$qt\sum_{a=1}^{E}h_a^2C_a+\sum_{i=1}^{N}r_i(\sum_{a=1}^{E}\frac{1}{2}h_a^2C_a\Delta_{ai}-u\sum_{a=1}^{E}h_a^2C_a)-su\sum_{a=1}^{E}h_a^2C_a=$$

$$=(\sum_{a=1}^{E}h_a^2C_a)(qt-u\sum_{i=1}^{N}r_i-su)+\sum_{a=1}^{E}\sum_{i=1}^{N}\frac{1}{2}r_ih_a^2C_a\Delta_{ai}=$$

$$=(\sum_{a=1}^{E}h_a^2C_a)(q\frac{1}{2}\frac{D-2-d}{D-2}-(D-2-q)\frac{1}{2}\frac{d}{D-2}+\frac{1}{2}(d-q))=0,$$

i.e. it is zero. The same is true for the terms containing χ_b ,

$$qu\sum_{b=1}^{M}v_b^2\chi_b-\sum_{i=1}^{N}r_i(\sum_{b=1}^{M}\frac{1}{2}v_b^2\chi_b\Lambda_{bi}-u\sum_{b=1}^{M}v_b^2\chi_b)-st\sum_{b=1}^{M}v_b^2\chi_b=$$

$$=(\sum_{b=1}^{M}v_b^2\chi_b)(qu+u\sum_{i=1}^{N}r_i-st)-\sum_{b=1}^{M}\sum_{i=1}^{N}\frac{1}{2}r_iv_b^2\chi_b\Lambda_{bi}=$$

$$=(\sum_{b=1}^{M}v_b^2\chi_b)(q\frac{1}{2}\frac{d}{D-2}+(D-2-q-s)\frac{1}{2}\frac{d}{D-2}-s\frac{D-2-d}{2(D-2)}-\frac{1}{2}(d-s))=0.$$

Non-diagonal part of the Einstein equations gives five types of relations. One type follows from the requirement of compensation $\partial_\alpha C_a\partial_\beta C_a$. It has the form

$$[-qt^2-\sum_{i=1}^{N}r_i(\frac{1}{4}\Delta_{ai}+u^2-u\Delta_{ai})-su^2]h_a^4=\frac{\alpha^2}{8}h_a^4-\frac{h_a^2}{2},$$

that is satisfied due to the requirement 4) and relation (2.29).

Another type of relations follows from the requirement of compensation $\partial_\alpha C_a\partial_\beta C_{a'}$, $a\neq a'$

$$\frac{\alpha^2}{8}+qt^2+\sum_{i=1}^{N}r_i(\frac{1}{4}\Delta_{ai}\Delta_{a'i}+u^2-u\Delta_{ai})+su^2=0,$$

that holds due to (1.10).

One more type is obtained from the requirement on the coefficient in front of $\partial_\alpha\chi_b\partial_\beta\chi_b$. It has the form

$$[-qu^2-\sum_{i=1}^{N}r_i(\frac{1}{4}\Lambda_{bi}+u^2-u\Lambda_{bi})-st^2]v_b^4=\frac{\alpha^2}{8}v_b^4-\frac{v_b^2}{2},$$

and takes place due to 4') and (2.35)

The next type of relations we get from the requirement of a compensation $\partial_\alpha\chi_b\partial_\beta\chi_b'$, $b\neq b'$

$$\frac{\alpha^2}{8}+qu^2+\sum_{i=1}^{N}r_i(\frac{1}{2}\Lambda_{bi}-u)(\frac{1}{2}\Lambda_{b'i}-u)+st^2=0,$$

that is equivalent to (1.13).

The last type of relations follows from the requirement of compensation $\partial_\alpha\chi_b\partial_\beta C_a$ and it looks like

$$\frac{\alpha^2}{8}-qut+\sum_{i=1}^{N}r_i(\frac{1}{2}\Lambda_{bi}-u)(\frac{1}{2}\Delta_{ai}-u)-stu=0. \tag{2.37}$$

Since the LHS of the above relations is

$$\frac{\alpha^2}{2}(D-2)-d^2+(D-2)\sum_{i=1}^{N}r_i\Lambda_{bi}\Delta_{ai}$$

so the requirement (2.37) is nothing but (1.14).

The form of the metric for this solution is

$$ds^2=(H_1^{-1}H_2^{-1}...H_E^{-1}U_1U_2...U_M)^{2t\sigma}(H_1H_2...H_E)^{\sigma}\times$$

$$\{(H_1H_2...H_EU_1U_2...U_M)^{-\sigma}\sum_{\mu,\nu=0}^{q-1}\eta_{\mu\nu}dy^\mu dy^\nu+$$

$$(U_1U_2...U_M)^{-\sigma}\sum_i(\prod_a H_a^{-\Delta_{ai}}\prod_b U_b^{\Lambda_{bi}})^{\sigma}\sum_{m_i=1}^{r_i}dz_i^{m_i}dz_i^{m_i}+\sum_\alpha dx^\alpha dx^\alpha\}, \tag{2.38}$$

where harmonic functions H_a and U_b are given by

$$H_a = 1 + \sum_c \frac{Q_{ac}}{|x - x_{ac}|^s}, \tag{2.39}$$

$$U_b = 1 + \sum_c \frac{P_{bc}}{|x - x_{bc}|^s}. \tag{2.40}$$

This representation of the metric is convenient for calculating the entropy. The hole area of the horizon is non-trivial and points x_{ac} and x_{bc} are not singularities when

1)all harmonic functions have the equal number of centers k and $x_{1c} = \ldots = x_{kc} = x_c$.

2) $2\sigma s(Eu + tM) = 2$

3) $\sigma s(\sum_a \Delta_{ai} - \sum_b \Lambda_{bi} + M) = 2$

4)$q = 1$ or

4')$q = 2$ and $s\sigma(E + M) - 2 = s$

Under these conditions for the case $q = 1$ the area of horizon has the form

$$\mathcal{A}_{D-2} = \omega_{s+1} L^{D-s-3} \sum_c (\prod_a Q_{ac} \prod_b P_{bc})^{\frac{\sigma}{2}}, \tag{2.41}$$

where ω_{s+1} is volume of $s+1$-dimensional sphere,

$$\omega_{s+1} = \frac{2\pi^{\frac{s+2}{2}}}{\Gamma\left(\frac{s+2}{2}\right)} \tag{2.42}$$

and L is a period of all y_n, $n = 1, .., D - s - 3$. To get non-trivial entropy for $q = 2$ one have to make a 'boost'

$$-dy_0^2 + dy_1^2 \longrightarrow dudv + K(x) du^2,$$

where

$$u = y_1 + y_0,\ v = y_1 - y_0,\ K(x) = 1 + \sum_c \frac{Q_c}{|x - x_c|^s},\ x_c = x_{ac}$$

In this case the area of horizon is given by

$$\mathcal{A}_{D-2} = \omega_{s+1} L^{D-s-3} \sum_c (\prod_a Q_{ac} \prod_b P_{bc})^{\frac{\sigma}{2}} Q_c^{\frac{1}{2}} \tag{2.43}$$

2.2 Examples

To describe particular examples of the general solution (1.12) it is convenient to present the incidence matrices Δ_{ai} and Λ_{bi} by special tables with E (or M) rows and $N+2$ columns (additional two columns correspond to subspaces of $\mathbf{M}$ with dimensions q and s, briefly q-column and s-column). If $\Delta_{ai}=1$ we put r_i symbols '$\circ$' in an appropriate cell of the table. If $\Lambda_{bi}=1$ we draw r_i symbols '$\bullet$' in a corresponding place of the table. Also we draw '$\circ$' and '$\bullet$' in q-columns and s-columns, respectively. We will not consider here the cases with $E+M=1$.

2.2.1 $D=11$, d-form and $\alpha=0$

First we will investigate the case $D=11$ and $\alpha=0$

1) *"electric" ansatz.*

In this case we have the following condition

$$q-\frac{d^2}{9}+\sum_{i=1}^{N} r_i\Delta_{ai}\Delta_{a'i}=0, \quad a\neq a' \tag{2.44}$$

where all values are integer. Therefore there are two types of solutions.

d=3, q=1, s=2

○	○ ○			
○		○ ○		
○			○ ○	

$$ds^2=(H_1H_2H_3)^{\frac{1}{3}}[-(H_1H_2H_3)^{-1}dy^2+H_1^{-1}(dy_1^2+dy_2^2)+H_2^{-1}(dy_3^2+dy_4^2)+$$
$$+H_3^{-1}(dy_5^2+dy_6^2)+dx^\alpha dx^\alpha] \tag{2.45}$$

For H_a in the form

$$H_a=1+\sum_c\frac{Q_{ac}}{|x-x_{ac}|} \tag{2.46}$$

we get non-zero area of the horizon if all positions of the poles x_{ac} coincide, $x_{ac}=x_c$,

$$\mathcal{A}_9=\omega_3L^6\sum_c(Q_{1c}Q_{2c}Q_{3c})^{1/2} \tag{2.47}$$

d=6, q=2, s=1

○ ○	○ ○	○ ○	
○ ○	○ ○		○ ○
○ ○		○ ○	○ ○

$$ds^2=(H_1H_2H_3)^{\frac{2}{3}}[(H_1H_2H_3)^{-1}(-dy^2+dy_1^2+K(x)du^2)+(H_1H_2)^{-1}(dy_2^2+dy_3^2)+$$
$$(H_1H_3)^{-1}(dy_4^2+dy_5^2)+(H_2H_3)^{-1}(dy_6^2+dy_7^2)+dx^\alpha dx^\alpha] \tag{2.48}$$

$$\mathcal{A}_9=4\pi L^7\sum_c(Q_cQ_{1c}Q_{2c}Q_{3c})^{1/2} \tag{2.49}$$

Note that the solutions presented in these tables correspond to "maximal" configuration in the sense that apart from the ansatzes presented by the above tables one can also consider ansatzes which can be obtained by deleting some rows of our matrices, but these solutions have zero area of the horizon.

The solution (2.45) has been discussed in [20, 37], the solution (2.48) was considered in [24].

2) *"magnetic" ansatz.*

In this case we have the following condition

$$s-\frac{d^2}{9}+\sum_{i=1}^{N} r_i\Lambda_{bi}\Lambda_{b'i}=0,\ b\neq b', \tag{2.50}$$

where all values are integer. Therefore there are similary two types of solutions.

d=3, q=2, s=1

			● ●	●
		● ●		●
	● ●			●

$$ds^2=(U_1U_2U_3)^{\frac{2}{3}}[(U_1U_2U_3)^{-1}(-dy^2+dy_1^2+K(x)du^2)+(U_1U_2)^{-1}(dy_2^2+dy_3^2)+$$

$$(U_1U_3)^{-1}(dy_4^2+dy_5^2)+(U_2U_3)^{-1}(dy_6^2+dy_7^2)+dx^\alpha dx^\alpha] \tag{2.51}$$

$$\mathcal{A}_9=4\pi L^7\sum_c(Q_cP_{1c}P_{2c}P_{3c})^{1/2} \tag{2.52}$$

d=6, q=1, s=2

		● ●	● ●	● ●
	● ●		● ●	● ●
	● ●	● ●		● ●

$$ds^2=(U_1U_2U_3)^{\frac{1}{3}}[-(U_1U_2U_3)^{-1}dy^2+U_1^{-1}(dy_1^2+dy_2^2)+U_2^{-1}(dy_3^2+dy_4^2)+$$

$$+U_3^{-1}(dy_5^2+dy_6^2)+dx^\alpha dx^\alpha] \tag{2.53}$$

$$\mathcal{A}_9=\omega_3L^6\sum_c(P_{1c}P_{2c}P_{3c})^{1/2} \tag{2.54}$$

These solutions are dual to the 'electric' ones.

3)*"electric" + "magnetic" ansatz.*

In this case we have the additional condition

$$-\frac{d^2}{9}+\sum_{i=1}^{N} r_i\Lambda_{bi}\Delta_{ai}=0 \tag{2.55}$$

Our strategy in finding solutions of (2.44), (2.50) and (2.55) is very simple. We take tables corresponding to electric and magnetic ansatzes and try to put on the corresponding configurations of the fields the additional condition (2.55). One can check that we cannot combine the maximal ansatz presented by (2.45) (or (2.48)) with maximal magnetic ansatz corresponding to (2.51) (or (2.53)), but we can combine a part of table (2.45) with a part of table (2.51) as well as a part of table (2.48) with a part of table (2.53) as it is presented in the tables connected with solutions (2.56) and (2.58), respectively. So, we have two types of solutions.

d=3, q=1, s=1

○	○	○				
○			○	○		
	●		●			●
		●		●		●

$$ds^2 = (H_1H_2)^{\frac{1}{3}}(U_1U_2)^{\frac{2}{3}}[-(H_1H_2U_1U_2)^{-1}dy^2 + (H_1U_2)^{-1}dy_1^2 + (H_1U_1)^{-1}dy_2^2 + \\ + (H_2U_2)^{-1}dy_3^2 + H_2{U_1}^{-1}dy_4^2 + (U_1U_2)^{-1}(dy_5^2 + dy_6^2 + dy_7^2) + dx^\alpha dx^\alpha] \tag{2.56}$$

$$\mathcal{A}_9 = 4\pi L^7 \sum_c (Q_{1c}Q_{2c}P_{1c}P_{2c})^{1/2} \tag{2.57}$$

d=6, q=1, s=1

○	○ ○ ○	○	○			
○	○ ○ ○			○	○	
	● ● ●	●		●		●
	● ● ●		●		●	●

$$ds^2 = (H_1H_2)^{\frac{2}{3}}(U_1U_2)^{\frac{1}{3}}[-(H_1H_2U_1U_2)^{-1}dy^2 + (H_1H_2)^{-1}(dy_1^2 + dy_2^2 + dy_3^2) + (H_1U_2)^{-1}dy_4^2 + \\ + (H_1U_1)^{-1}dy_5^2 + (H_2U_2)^{-1}dy_6^2 + (H_2U_1)^{-1}dy_7^2 + dx^\alpha dx^\alpha] \tag{2.58}$$

$$\mathcal{A}_9 = 4\pi L^7 \sum_c (Q_{1c}Q_{2c}P_{1c}P_{2c})^{1/2} \tag{2.59}$$

The solution (2.56) has been constructed in [24] and it is dual to (2.58).

2.2.2 D=10, d-form and $\alpha \neq 0$.

1)*"electric" ansatz.*

In this case we have the following condition

$$\frac{\alpha^2}{2} + q - \frac{d^2}{9} + \sum_{i=1}^{N} r_i \Delta_{ai}\Delta_{a'i} = 0,\ a \neq a',$$

where all values are integer. Therefore there are four types of solutions.

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1

○	○ ○			
○		○ ○		
○			○ ○	

$$ds^2 = (H_1H_2H_3)^{\frac{3}{8}}[-(H_1H_2H_3)^{-1}dy^2 + H_1^{-1}(dy_1^2 + dy_2^2) + H_2^{-1}(dy_3^2 + dy_4^2) +$$

$$+ H_3^{-1}(dy_5^2 + dy_6^2) + dx^\alpha dx^\alpha] \tag{2.60}$$

All centers of harmonic functions are points of singularity.

d=4,$\alpha = \pm\sqrt{2}$, q=1, s=1

○	○	○	○				
○				○	○	○	

$$ds^2 = (H_1H_2)^{\frac{1}{3}}[-(H_1H_2)^{-\frac{2}{3}}dy^2 + H_1^{-\frac{2}{3}}(dy_1^2 + dy_2^2 + dy_3^2) +$$

$$+ H_2^{-\frac{2}{3}}(dy_4^2 + dy_5^2 + dy_6^2) + dx^\alpha dx^\alpha] \tag{2.61}$$

The area of horizon is equal to zero.

d=4, $\alpha = 0$, q=1, s=1

○	○	○	○				
○	○			○	○		
○		○		○		○	
○			○		○	○	

$$ds^2 = (H_1H_2H_3H_4)^{\frac{1}{2}}[-(H_1H_2H_3H_4)^{-1}dy^2 + (H_1H_2)^{-1}dy_1^2 + (H_1H_3)^{-1}dy_2^2 + (H_1H_4)^{-1}dy_3^2 +$$

$$+ (H_2H_3)^{-1}dy_4^2 + (H_2H_4)^{-1}dy_5^2 + (H_3H_4)^{-1}dy_6^2 + dx^\alpha dx^\alpha] \tag{2.62}$$

$$\mathcal{A}_8 = 4\pi L^6 \sum_c (Q_{1c}Q_{2c}Q_{3c}Q_{4c})^{1/2} \tag{2.63}$$

This solution was discussed in [30].

d=5, $\alpha = \pm\frac{1}{2}$, q=1,s=1

○	○	○	○	○			
○	○	○			○	○	
○			○	○	○	○	

$$ds^2 = (H_1H_2H_3)^{\frac{5}{8}}[-(H_1H_2H_3)^{-1}dy^2 + (H_1H_2)^{-1}(dy_1^2 + dy_2^2) + (H_1H_3)^{-1}(dy_3^2 + dy_4^2) +$$

$$+ (H_2H_3)^{-1}(dy_5^2 + dy_6^2) + dx^\alpha dx^\alpha] \tag{2.64}$$

The area of horizon is equal to zero.

2) *"magnetic" ansatz.*

In this case we have the following condition

$$\frac{\alpha^2}{2} + s - \frac{d^2}{9} + \sum_{i=1}^{N} r_i \Lambda_{bi}\Lambda_{b'i} = 0 \;\; b \neq b'$$

where all values are integer. Therefore there are also four types of solutions.

d=3, $\alpha=\pm\frac{1}{2}$, q=1, s=1

			● ●	●
		● ●		●
	● ●			●

$$ds^2=(U_1U_2U_3)^{\frac{5}{8}}[-(U_1U_2U_3)^{-1}dy^2+(U_1U_2)^{-1}(dy_1^2+dy_2^2)+(U_1U_3)^{-1}(dy_3^2+dy_4^2)+$$
$$+(U_2U_3)^{-1}(dy_5^2+dy_6^2)+dx^\alpha dx^\alpha] \qquad (2.65)$$

The entropy is equal to zero.

d=4, $\alpha=\pm\sqrt{2}$, q=1, s=1

		● ● ●	●
	● ● ●		●

$$ds^2=(U_1U_2)^{\frac{1}{3}}[-(U_1U_2)^{-\frac{2}{3}}dy^2+U_1^{-\frac{2}{3}}(dy_1^2+dy_2^2+dy_3^2)+$$
$$+U_2^{-\frac{2}{3}}(dy_4^2+dy_5^2+dy_6^2)+dx^\alpha dx^\alpha] \qquad (2.66)$$

The area of horizon is equal to zero.

d=4, $\alpha=0$, q=1, s=1

				●	●	●	●
		●	●			●	●
	●		●		●		●
	●	●		●			●

$$ds^2=(U_1U_2U_3U_4)^{\frac{1}{2}}[-(U_1U_2U_3U_4)^{-1}dy^2+(U_1U_2)^{-1}dy_1^2+(U_1U_3)^{-1}dy_2^2+(U_1U_4)^{-1}dy_3^2+$$
$$+(U_2U_3)^{-1}dy_4^2+(U_2U_4)^{-1}dy_5^2+(U_3U_4)^{-1}dy_6^2+dx^\alpha dx^\alpha] \qquad (2.67)$$

$$\mathcal{A}_8=4\pi L^6\sum_c(Q_{1c}Q_{2c}Q_{3c}Q_{4c})^{1/2} \qquad (2.68)$$

d=5, $\alpha=\pm\frac{1}{2}$, q=1, s=1

		● ●	● ●	●
	● ●		● ●	●
	● ●	● ●		●

$$ds^2=(U_1U_2U_3)^{\frac{3}{8}}[-(U_1U_2U_3)^{-1}dy^2+U_1^{-1}(dy_1^2+dy_2^2)+U_2^{-1}(dy_3^2+dy_4^2)+$$
$$+U_3^{-1}(dy_5^2+dy_6^2)+dx^\alpha dx^\alpha] \qquad (2.69)$$

This metric has singularities in the centers of U_b

The solutions (2.65),(2.66),(2.67) and (2.69) are dual to (2.64),(2.61),(2.62) and (2.60) respectively.

3) *"electric" + "magnetic" ansatz.* In this case we have the additional condition

$$\frac{\alpha^2}{2}-\frac{d^2}{9}+\sum_{i=1}^N r_i\Lambda_{bi}\Delta_{ai}=0$$

Again we have three types of solutions.

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1

○	○	○				
○			○	○		
	●		●			●
		●		●		●

$$ds^2 = (H_1H_2)^{\frac{3}{8}}(U_1U_2)^{\frac{5}{8}}[-(H_1H_2U_1U_2)^{-1}dy^2+(H_1U_2)^{-1}dy_1^2+(H_1U_1)^{-1}dy_2^2+(H_2U_2)^{-1}dy_3^2+$$
$$+ (H_2U_1)^{-1}dy_4^2 + (U_1U_2)^{-1}(dy_5^2 + dy_6^2) + dx^\alpha dx^\alpha] \qquad (2.70)$$

$$\mathcal{A}_8 = 4\pi L^6 \sum_c (Q_{1c}Q_{2c}P_{1c}P_{2c})^{1/2} \qquad (2.71)$$

$d = 4,\ \alpha = \pm\sqrt{2},\ q = 3$

○	○ ○	○		
		●	● ●	●

$$ds^2 = (HU)^{\frac{1}{3}}[(HU)^{-\frac{2}{3}}(-dy^2 + dy_1^2 + dy_2^2)+$$
$$+ H^{-\frac{2}{3}}dy_3^2 + U^{-\frac{2}{3}}dy_4^2 + dx^\alpha dx^\alpha] \qquad (2.72)$$

In this case $q = 3$, therefore the entropy is zero.

d=5, $\alpha = \pm\frac{1}{2}$, q=1, s=1

○	○ ○	○	○			
○	○ ○			○	○	
	● ●	●		●		●
	● ●		●		●	●

$$ds^2 = (H_1H_2)^{\frac{5}{8}}(U_1U_2)^{\frac{3}{8}}[-(H_1H_2U_1U_2)^{-1}dy^2 + (H_1H_2)^{-1}(dy_1^2 + dy_2^2) + (H_1U_2)^{-1}dy_3^2+$$
$$+ (H_1U_1)^{-1}dy_4^2 + (H_2U_2)^{-1}dy_5^2 + (H_2U_1)^{-1}dy_6^2 + dx^\alpha dx^\alpha] \qquad (2.73)$$

$$\mathcal{A}_8 = 4\pi L^6 \sum_c (P_{1c}P_{2c}Q_{1c}Q_{2c})^{1/2} \qquad (2.74)$$

2.2.3 D=18, d-form, and $\alpha = 0$

Let us demonstrate our method in the complicated case $D = 18$. For simplicity we consider only the 'electric' ansatz.

$d = 4,\ \alpha = 0,\ q = 1,\ s = 3$

○	○ ○ ○				
○		○ ○ ○			
○			○ ○ ○		
○				○ ○ ○	

$$ds^2 = (H_1H_2H_3H_4)^{\frac{1}{6}}[-(H_1H_2H_3H_4)^{-\frac{2}{3}}dy^2 + (H_1)^{-\frac{2}{3}}(dy_1^2 + dy_2^2 + dy_3^2)+$$
$$+(H_2)^{-\frac{2}{3}}(dy_4^2 + dy_5^2 + dy_6^2) + (H_3)^{-\frac{2}{3}}(dy_7^2 + dy_8^2 + dy_9^2)+$$
$$+ (H_4)^{-\frac{2}{3}}(dy_{10}^2 + dy_{11}^2 + dy_{12}^2) + dx^\alpha dx^\alpha] \qquad (2.75)$$

$$\mathcal{A}_{16} = \omega_4 L^{12} \sum_c (Q_{1c}Q_{2c}Q_{3c}Q_{4c})^{1/3} \tag{2.76}$$

d=8, $\alpha = 0$, q=1, s=1

○	○	○	○	○	○	○	○								
○	○	○	○					○	○	○	○				
○	○	○		○				○				○	○	○	
○	○				○	○			○	○		○	○		
○			○		○		○	○		○		○		○	
○				○		○	○	○		○	○		○		
○		○			○		○		○		○		○	○	
○			○	○		○			○		○	○		○	

$$ds^2 = (H_1H_2H_3H_4H_5H_6H_7H_8)^{\frac{1}{4}}[-(H_1H_2H_3H_4H_5H_6H_7H_8)^{-\frac{1}{2}}dy^2 + (H_1H_2H_3H_4)^{-\frac{1}{2}}dy_1^2 +$$

$$+(H_1H_2H_3H_7)^{-\frac{1}{2}}dy_2^2 + +(H_1H_2H_5H_8)^{-\frac{1}{2}}dy_3^2 + (H_1H_3H_6H_8)^{-\frac{1}{2}}dy_4^2 +$$

$$+(H_1H_4H_5H_7)^{-\frac{1}{2}}dy_5^2 + +(H_1H_4H_6H_8)^{-\frac{1}{2}}dy_6^2 + (H_1H_5H_6H_7)^{-\frac{1}{2}}dy_7^2 + (H_2H_3H_5H_6)^{-\frac{1}{2}}dy_8^2 +$$

$$+(H_2H_4H_7H_8)^{-\frac{1}{2}}dy_9^2 + (H_2H_4H_5H_6)^{-\frac{1}{2}}dy_{10}^2 + (H_2H_6H_7H_8)^{-\frac{1}{2}}dy_{11}^2 + (H_3H_4H_5H_8)^{-\frac{1}{2}}dy_{12}^2 +$$

$$(H_3H_4H_6H_7)^{-\frac{1}{2}}dy_{13}^2 + (H_3H_5H_7H_8)^{-\frac{1}{2}}dy_{14}^2 + dx^\alpha dx^\alpha] \tag{2.77}$$

$$\mathcal{A}_{16} = 4\pi L^{14} \sum_c (Q_{1c}Q_{2c}Q_{3c}Q_{4c}Q_{5c}Q_{6c}Q_{7c}Q_{8c})^{1/4} \tag{2.78}$$

d=12, $\alpha = 0$, q=3, s=1

○ ○ ○	○ ○ ○	○ ○ ○	○ ○ ○		
○ ○ ○	○ ○ ○	○ ○ ○		○ ○ ○	
○ ○ ○	○ ○ ○		○ ○ ○	○ ○ ○	
○ ○ ○		○ ○ ○	○ ○ ○	○ ○ ○	

$$ds^2 = (H_1H_2H_3H_4)^{\frac{1}{2}}[(H_1H_2H_3H_4)^{-\frac{2}{3}}(-dy^2 + dy_1^2 + dy_2^2) + (H_1H_2H_3)^{-\frac{2}{3}}(dy_3^2 + dy_4^2 + dy_5^2) +$$

$$+(H_1H_2H_4)^{-\frac{2}{3}}(dy_6^2 + dy_7^2 + dy_8^2) + (H_1H_3H_4)^{-\frac{2}{3}}(dy_9^2 + dy_{10}^2 + dy_{11}^2) +$$

$$+ (H_2H_3H_4)^{-\frac{2}{3}}(dy_{12}^2 + dy_{13}^2 + dy_{14}^2) + dx^\alpha dx^\alpha] \tag{2.79}$$

In this solution $q = 3$, the area of horizon is zero.

3 Gravity coupled with d_1-form, d_2-form and dilaton

In this section we will consider soliton solution of the form (1.3) for the theory

$$I=\frac{1}{2\kappa^2}\int d^Dx\sqrt{-g}(R-\frac{1}{2}(\nabla\phi)^2-\frac{e^{-\alpha\phi}}{2(d_1+1)!}F^2_{d_1+1}-\frac{e^{-\beta\phi}}{2(d_2+1)!}F^2_{d_2+1}), \tag{3.1}$$

We will follow the same strategy of finding solutions as before. For simplicity let us present the main steps of calculations only for pure 'electric' field but write only the answer for a general 'electric' + 'magnetic' ansatz.

For d_1 and d_2-form we are going to consider a class of ansatzes corresponding to E_1 and E_2 electric charges. Each ansatz we will connect with two incidence matrices $\Delta^{(1)}_{ai}$, $a=1..E_1$, $i=1..N$, $\Delta^{(2)}_{bi}$, $b=1..E_2$, $i=1..N$, which apart the properties 1), 2), 3) and 5) mentioned in the previous section satisfy the following conditions

4) $\sum_{i=1}^N r_i\Delta^{(1)}_{ai}=d_1-q$ for all a;

$\sum_{i=1}^N r_i\Delta^{(2)}_{ai}=d_2-q$ for all a;

Electric fields are assumed in the form

$$\mathcal{A}^{(1)}=\sum_{a=1}^{E_1}h^{(1)}_a e^{C_a(x)^{(1)}}\omega_0^q\prod_{i|\Delta^{(1)}_{ai}=1}\wedge\,\omega_i \tag{3.2}$$

$$\mathcal{A}^{(2)}=\sum_{a=1}^{E_2}h^{(2)}_a e^{C_a(x)^{(2)}}\omega_0^q\prod_{i|\Delta^{(2)}_{ai}=1}\wedge\,\omega_i \tag{3.3}$$

Components of the energy-momentum tensor for d_1-form and d_2-form are very similar except that in the exponent corresponding to the d_2-field new parameter β enters instead of α .

"No-force" conditions in this case have the form

$$-\alpha\phi-2qA-2\sum_{i=1}^N\Delta^{(1)}_{ai}r_iF_i+2C^{(1)}_a=0,\ i=1..E_1 \tag{3.4}$$

$$-\beta\phi-2qA-2\sum_{i=1}^N\Delta^{(2)}_{ai}r_iF_i+2C^{(2)}_a=0,\ i=1..E_2 \tag{3.5}$$

From Maxwell's equations under conditions (3.4) and (3.5) we get

$$\Delta C^{(1)}_a=(\partial C^{(1)}_a)^2,\quad \Delta C^{(2)}_a=(\partial C^{(2)}_a)^2$$

Having in our disposal these expressions the equation of motion for the dilaton as well as the Einstein equations (apart from the non-diagonal part for γ-components can be solved if one assume that

$$A=\sum_{a=1}^{E_1}t^{(1)}{h^{(1)}_a}^2C^{(1)}_a+\sum_{a=1}^{E_2}t^{(2)}{h^{(2)}_a}^2C^{(2)}_a \tag{3.6}$$

$$F_i=\sum_{a=1}^{E_1}{h^{(1)}_a}^2C^{(1)}_a(\frac{1}{2}\Delta^{(1)}_{ai}-u^{(1)})+\sum_{a=1}^{E_2}{h^{(2)}_a}^2C^{(2)}_a(\frac{1}{2}\Delta^{(2)}_{ai}-u^{(2)}) \tag{3.7}$$

$$B = -u^{(1)} \sum_{a=1}^{E_1} {h_a^{(1)}}^2 C_a^{(1)} - u^{(2)} \sum_{a=1}^{E_2} {h_a^{(2)}}^2 C_a^{(2)} \tag{3.8}$$

$$\phi = \frac{\alpha}{2} \sum_{a=1}^{E_1} {h_a^{(1)}}^2 C_a^{(1)} + \frac{\beta}{2} \sum_{a=1}^{E_2} {h_a^{(2)}}^2 C_a^{(2)} \tag{3.9}$$

where $t^{(1)},\ t^{(2)},\ u^{(1)}$ and $u^{(2)}$ are given by

$$t^{(1)} = \frac{D-2-d_1}{2(D-2)} \tag{3.10}$$

$$t^{(2)} = \frac{D-2-d_2}{2(D-2)} \tag{3.11}$$

$$u^{(1)} = \frac{d_1}{2(D-2)} \tag{3.12}$$

$$u^{(2)} = \frac{d_2}{2(D-2)} \tag{3.13}$$

Having the formulae (3.6), (3.7) and (3.8) one can immediate verify that the relation (2.3) holds.

In the same way as it has been done in Section 2 one can find that consistency conditions of equations (3.6), (3.7), 3.8) and (3.9) with (4.39) and (4.40) force us to put the following restrictions

$$(D-2)(\frac{\alpha^2}{2} + q) - d_1^2 + (D-2) \sum_{i=1}^{N} r_i \Delta_{ai}^{(1)} \Delta_{a'i}^{(1)} = 0, \quad a \neq a' \tag{3.14}$$

$$(\frac{\alpha\beta}{2} + q)(D-2) - d_1 d_2 + (D-2) \sum_{i=1}^{N} r_i \Delta_{ai}^{(1)} \Delta_{a'i}^{(2)} = 0 \tag{3.15}$$

$$(D-2)(\frac{\beta^2}{2} + q) - d_2^2 + (D-2) \sum_{i=1}^{N} r_i \Delta_{ai}^{(2)} \Delta_{a'i}^{(2)} = 0 \quad a \neq a' \tag{3.16}$$

and take h_a in the form

$${h_a^{(1)}}^2 = \sigma^{(1)} = \frac{1}{t^{(1)} d_1 + \frac{\alpha^2}{4}}, \quad {h_a^{(2)}}^2 = \sigma^{(2)} = \frac{1}{t^{(2)} d_2 + \frac{\beta^2}{4}} \tag{3.17}$$

One can check that if these conditions are satisfied then the rest of the Einstein equations are hold. Introducing the notations $\alpha = \alpha^{(1)},\ \beta = \alpha^{(2)}$ we can rewrite (3.14) –(3.16) as

$$(1 - \delta_{IJ}\delta_{aa'})\{\frac{\alpha^{(I)}\alpha^{(J)}}{2} + q - \frac{d_I d_J}{D-2} + \sum_{i=1}^{N} r_i \Delta_{ai}^{(I)} \Delta_{a'i}^{(J)}\} = 0,\ I,\ J = 1,\ 2 \tag{3.18}$$

Let us present the metric corresponding to this solution

$$ds^2 = (H_1^{(1)} H_2^{(1)} ... H_{E_1}^{(1)})^{2u^{(1)}\sigma^{(1)}} (H_1^{(2)} H_2^{(2)} ... H_{E_2}^{(2)})^{2u^{(2)}\sigma^{(2)}}$$

$$\{(H_1^{(1)}H_2^{(1)}...H_{E_1}^{(1)})^{-\sigma^{(1)}}(H_1^{(2)}H_2^{(2)}...H_{E_2}^{(2)})^{-\sigma^{(2)}}\eta_{\mu\nu}dy^\mu dy^\nu+$$

$$+\sum_i(\prod_a H_a^{(1)^{-\Delta_{ai}^{(1)}}})^{\sigma^{(1)}}(\prod_a H_a^{(2)^{-\Delta_{ai}^{(2)}}})^{\sigma^{(2)}}\sum_{m_i=1}^{r_i}dz^{m_i}dz^{m_i}+\sum_\alpha dx^\alpha dx^\alpha], \qquad (3.19)$$

In the most general case with electric and magnetic fields we have additional restrictions on the incidence matrices

$$(1-\delta_{IJ}\delta_{bb'})\{\frac{\alpha^{(I)}\alpha^{(J)}}{2}+s-\frac{d_Id_J}{D-2}+\sum_{i=1}^N r_i\Lambda_{bi}^{(I)}\Lambda_{b'i}^{(J)}\}=0, \qquad (3.20)$$

$$\frac{\alpha^{(I)}\alpha^{(J)}}{2}+s-\frac{d_Id_J}{D-2}+\sum_{i=1}^N r_i\Delta_{ai}^{(I)}\Lambda_{b'i}^{(J)}=0,\ I,\ J=1,\ 2 \qquad (3.21)$$

and

$$v_m^{(I)^2}=v^{(I)^2}=\frac{1}{t^{(I)}d_I+\frac{\alpha_I^2}{4}},\ I=1,\ 2 \qquad (3.22)$$

The metric has the form

$$ds^2=(H_1^{(1)}H_2^{(1)}...H_{E_1}^{(1)})^{2u^{(1)}\sigma^{(1)}}(U_1^{(1)}U_2^{(1)}...U_{M_1}^{(1)})^{2t^{(1)}\sigma^{(1)}}(H_1^{(2)}H_2^{(2)}...H_{E_2}^{(2)})^{2u^{(2)}\sigma^{(2)}}$$

$$(U_1^{(2)}U_2^{(2)}...U_{M_2}^{(2)})^{2t^{(2)}\sigma^{(2)}}\{(H_1^{(1)}H_2^{(1)}...H_{E_1}^{(1)}U_1^{(1)}U_2^{(1)}...U_{M_1}^{(1)})^{-\sigma^{(1)}}(H_1^{(2)}H_2^{(2)}...H_{E_2}^{(2)}$$

$$U_1^{(2)}U_2^{(2)}...U_{M_2}^{(2)})^{-\sigma^{(2)}}\eta_{\mu\nu}dy^\mu dy^\nu+\sum_i(\prod_a H_a^{(1)^{-\Delta_{ai}^{(1)}}}\prod_b U_b^{(1)^{\Lambda_{bi}^{(1)}}})^{\sigma^{(1)}}$$

$$(\prod_a H_a^{(2)^{-\Delta_{ai}^{(2)}}}\prod_b U_b^{(2)^{\Lambda_{ai}^{(2)}}})^{\sigma^{(2)}}\sum_{m_i=1}^{r_i}dz^{m_i}dz^{m_i}+\sum_\alpha dx^\alpha dx^\alpha], \qquad (3.23)$$

Note that in the case of one dilaton field and an arbitrary number of d_I-forms, $I=1,..,S$, requirements on $2S$ incidence matrices are also given by (3.18), (3.20) and (3.21) with $I,\ J=1,..,S$ and corresponding metric is an obvious generalization of (3.23). Furthermore, the same method could be applied to the case with more then one dilaton fields.

3.1 Examples

3.1.1 D=11

We present below only solutions with at list two branches fot both fields. There are more solutions with one branch for earch field.

1) *"electric" ansatzes.*

In this case we get

$$q-\frac{d_1d_2}{9}+\sum_{i=1}^N r_i\Delta_{ai}^1\Delta_{a'i}^2=0$$

and we have two types of solution

$d_1=3$, q=1, s=1

$d_2=6$, q=2, s=1



$$ds^2=(H^{(1)}_1H^{(1)}_2)^{\frac{1}{3}}(H^{(2)}_1H^{(2)}_2)^{\frac{2}{3}}[-(H^{(1)}_1H^{(1)}_2H^{(2)}_1H^{(2)}_2)^{-1}dy^2+$$
$$(H^{(1)}_1H^{(2)}_1)^{-1}dy_1^2+(H^{(1)}_1H^{(2)}_2)^{-1}dy_2^2+(H^{(1)}_2H^{(2)}_1)^{-1}dy_3^2+(H^{(1)}_2H^{(2)}_2)^{-1}dy_4^2+$$
$$(H^{(2)}_1H^{(2)}_2)^{-1}(dy_5^2+dy_6^2+dy_7^2)+dx^\alpha dx^\alpha] \tag{3.24}$$

$$\mathcal{A}_9=4\pi L^7\sum_c(Q^{(1)}_{1c}Q^{(1)}_{2c}Q^{(2)}_{1c}Q^{(2)}_{2c})^{1/2} \tag{3.25}$$

2) *"magnetic" ansatzes*
In this case we get

$$s-\frac{d_1d_2}{9}+\sum_{i=1}^N r_i\Lambda^1_{bi}\Lambda^2_{b'i}=0$$

and corresponding solutions are

$d_1=3$, q=1, s=1			●		●		●
		●		●			●
$d_2=6$, q=1, s=1				●	●	● ● ●	●
		●	●			● ● ●	●

$$ds^2=(U^{(1)}_1U^{(1)}_2)^{\frac{2}{3}}(U^{(2)}_1U^{(2)}_2)^{\frac{1}{3}}[-U^{(1)}_1U^{(1)}_2U^{(2)}_1U^{(2)}_2dy^2+$$
$$+(U^{(1)}_1U^{(2)}_1)^{-1}dy_1^2+(U^{(1)}_2U^{(2)}_1)^{-1}dy_2^2+$$
$$+(U^{(1)}_1U^{(2)}_2)^{-1}dy_3^2+(U^{(1)}_2U^{(2)}_2)^{-1}dy_4^2+(U^{(1)}_1U^{(1)}_2)^{-1}(dy_5^2+dy_6^2+dy_7^2)+dx^\alpha dx^\alpha] \tag{3.26}$$

$$\mathcal{A}_9=4\pi L^7\sum_c(P^{(1)}_{1c}P^{(1)}_{2c}P^{(2)}_{1c}P^{(2)}_{2c})^{1/2} \tag{3.27}$$

3)*"electric" and "magnetic" ansatzes.*
New conditions are

$$-\frac{d_1d_2}{9}+\sum_{i=1}^N r_i\Lambda^1_{bi}\Delta^2_{ai}=0$$
$$-\frac{d_1d_2}{9}+\sum_{i=1}^N r_i\Lambda^2_{bi}\Delta^1_{ai}=0$$

We obtained three classes of solutions

$d_1=3$, q=1, s=1	○	○	○				
	○			○	○		
			●		●		●
$d_2=6$, q=1,s=1	○		○		○	○ ○ ○	

$$ds^2=(H^{(1)}_1H^{(1)}_2)^{\frac{1}{3}}(UH^{(2)})^{\frac{2}{3}}[-(H^{(1)}_1H^{(1)}_2UH^{(2)})^{-1}dy^2+(H^{(1)}_1U)^{-1}dy_1^2+(H^{(1)}_1H^{(2)})^{-1}dy_2^2+$$

$$+ (H_2^{(1)}U)^{-1}dy_3^2 + (H_2^{(1)}H^{(2)})^{-1}dy_4^2 + (UH^{(2)})^{-1}(dy_5^2 + dy_6^2 + dy_7^2) + dx^\alpha dx^\alpha] \qquad (3.28)$$

$d_1 = 6$, q=1, s=1

$d_2 = 3$, q=1, s=1



$$ds^2 = (H_1^{(1)}H_2^{(1)})^{\frac{2}{3}}(UH^{(2)})^{\frac{1}{3}}[-(H_1^{(1)}H_2^{(1)}UH^{(2)})^{-1}dy^2 +$$
$$+ (H_1^{(1)}H_2^{(1)})^{-1}(dy_1^2 + dy_2^2 + dy_3^2) + (H_1^{(1)}U)^{-1}dy_4^2 +$$
$$+ (H_1^{(1)}H^{(2)})^{-1}dy_5^2 + (H_2^{(1)}U)^{-1}dy_6^2 + (H_2^{(1)}H^{(2)})^{-1}dy_7^2 + dx^\alpha dx^\alpha] \qquad (3.29)$$

$d_1 = 3$, q=1, s=1

$d_2 = 6$, q=1, s=1



$$ds^2 = (H^{(1)}U^{(2)})^{\frac{1}{3}}(H^{(2)}U^{(1)})^{\frac{2}{3}}[-(H^{(1)}U^{(1)}H^{(2)}U^{(2)})^{-1}dy^2 +$$
$$+ (H^{(1)}U^{(1)})^{-1}dy_1^2 + (H^{(1)}H^{(2)})^{-1}dy_2^2 +$$
$$+ (U^{(1)}U^{(2)})^{-1}dy_3^2 + (H^{(2)}U^{(2)})^{-1}dy_4^2 + (H^{(2)}U^{(1)})^{-1}(dy_5^2 + dy_6^2 + dy_7^2) + dx^\alpha dx^\alpha] \qquad (3.30)$$

3.1.2 D=10

Now we will investigate the case $D = 10$ with two fields.

1) *"electric" ansatz.*

In this case we have the following condition

$$\frac{\alpha\beta}{2} + q - \frac{d_1 d_2}{8} + \sum_{i=1}^{N} r_i \Delta_{ai}^{(1)}\Delta_{a'i}^{(2)} = 0$$

where all values are integer. Therefore there are following types of solutions.

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1

d=5, $\beta = \mp\frac{1}{2}$, q=1, s=1



$$ds^2 = (H_1^{(1)}H_2^{(1)})^{\frac{3}{8}}(H_1^{(2)}H_2^{(2)})^{\frac{5}{8}}[-(H_1^{(1)}H_2^{(1)}H_1^{(2)}H_2^{(2)})^{-1}dy^2 + (H_1^{(1)}H_1^{(2)})^{-1}dy_1^2 +$$
$$+ (H_1^{(1)}H_2^{(2)})^{-1}dy_2^2 + + (H_2^{(1)}H_1^{(2)})^{-1}dy_3^2 + (H_2^{(1)}H_2^{(2)})^{-1}dy_4^2 +$$

$$+ (H_1^{(2)}H_2^{(2)})^{-1}(dy_5^2 + dy_6^2) + dx^\alpha dx^\alpha] \tag{3.31}$$

$$\mathcal{A}_8 = 4\pi L^6 \sum_c (Q_{1c}^{(1)} Q_{2c}^{(1)} Q_{1c}^{(2)} Q_{2c}^{(2)})^{1/2} \tag{3.32}$$

There are many solutions with two fieldes when each of them has only one electric or magnetic component. For example

$d_1 = 3,\ \alpha = \pm\frac{\sqrt{7}}{2}$

$d_2 = 3,\ \beta = \mp\frac{\sqrt{7}}{2}$



$$ds^2 = (H_1H_2)^{\frac{3}{11}}[(H_1H_2)^{-\frac{8}{11}}(-dy^2 + dy_1^2) + H_1^{-\frac{8}{11}}dy_2^2 + H_2^{-\frac{8}{11}}dy_3^2 + dx^\alpha dx^\alpha] \tag{3.33}$$

Centers of harmonic functions are singularities. A family of such solutions was discussed in [31]

2) *"magnetic" ansatz.* In this case we have the following condition

$$\frac{\alpha\beta}{2} + s - \frac{d_1d_2}{8} + \sum_{i=1}^{N} r_i \Lambda_{bi}^{(1)} \Lambda_{b'i}^{(2)} = 0$$

where all values are integer. Therefore there are three types of solutions.

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1

d=5, $\beta = \mp\frac{1}{2}$, q=1, s=1



$$ds^2 = (U_1^{(1)}U_2^{(1)})^{\frac{5}{8}}(U_1^{(2)}U_2^{(2)})^{\frac{3}{8}}[-(U_1^{(1)}U_2^{(1)}U_1^{(2)}U_2^{(2)})^{-1}dy^2 + (U_1^{(1)}U_2^{(1)})^{-1}(dy_1^2 +$$
$$+ dy_2^2) + (U_1^{(1)}U_1^{(2)})^{-1}dy_3^2 + (U_1^{(1)}U_2^{(2)})^{-1}dy_4^2 +$$
$$+ (U_2^{(1)}U_1^{(2)})^{-1}dy_5^2 + (U_2^{(1)}U_2^{(2)})^{-1}dy_6^2 + dx^\alpha dx^\alpha] \tag{3.34}$$

$$\mathcal{A}_8 = 4\pi L^6 \sum_c (P_{1c}^{(1)} P_{2c}^{(1)} P_{1c}^{(2)} P_{2c}^{(2)})^{1/2} \tag{3.35}$$

3)*"electric" + "magnetic"* ansatz. In this case we have the additional conditions

$$\frac{\alpha\beta}{2} - \frac{d_1d_2}{8} + \sum_{i=1}^{N} r_i \Lambda_{bi}^{(1)} \Delta_{ai}^{(2)} = 0,\ \frac{\alpha\beta}{2} - \frac{d_1d_2}{8} + \sum_{i=1}^{N} r_i \Lambda_{bi}^{(2)} \Delta_{ai}^{(1)} = 0$$

We have three types of solutions.

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1

d=5, $\beta = \mp\frac{1}{2}$, q=1, s=1



$$ds^2 = (H_1^{(1)} H_2^{(1)})^{\frac{3}{8}} (UH^{(2)})^{\frac{5}{8}} [-(H_1^{(1)} H_2^{(1)} UH^{(2)})^{-1} dy^2 + (H^{(2)}U)^{-1}(dy_1^2 + dy_2^2) +$$
$$+ (H_1^{(1)} U)^{-1} dy_3^2 + (H_2^{(1)} U)^{-1} dy_4^2 +$$
$$+ (H_1^{(1)} H^{(2)})^{-1} dy_5^2 + (H_2^{(1)} H^{(2)})^{-1} dy_6^2 + dx^\alpha dx^\alpha] \qquad (3.36)$$

d=5, $\alpha = \pm\frac{1}{2}$, q=1, s=1	○	○	○	○	○			
	○	○	○			○	○	
		●	●	●		●		●
d=3, $\beta = \mp\frac{1}{2}$, q=1, s=1	○			○		○		

$$ds^2 = (H_1^{(1)} H_2^{(1)})^{\frac{5}{8}} (UH^{(2)})^{\frac{3}{8}} [-(H_1^{(1)} H_2^{(1)} UH^{(2)})^{-1} dy^2 + (H_1^{(1)} H_2^{(1)})^{-1}(dy_1^2 +$$
$$+ dy_2^2) + (H_1^{(1)} H^{(2)})^{-1} dy_3^2 + (H_1^{(1)} U)^{-1} dy_4^2 +$$
$$+ (H_2^{(1)} H^{(2)})^{-1} dy_5^2 + (H_2^{(1)} U)^{-1} dy_6^2 + dx^\alpha dx^\alpha] \qquad (3.37)$$

d=3, $\alpha = \pm\frac{1}{2}$, q=1, s=1	○			○		○		
				●	●			●
d=5, $\beta = \mp\frac{1}{2}$, q=1, s=1	○	○	○	○	○			
		●	●	●		●		●

$$ds^2 = (H^{(1)} U^{(2)})^{\frac{3}{8}} (H^{(2)} U^{(1)})^{\frac{5}{8}} [-(H^{(1)} U^{(1)} H^{(2)} U^{(2)})^{-1} dy^2 + (H^{(2)} U^{(1)})^{-1}(dy_1^2 + dy_2^2) +$$
$$+ (H^{(1)} H^{(2)})^{-1} dy_3^2 + (H^{(2)} U^{(2)})^{-1} dy_4^2 +$$
$$+ (H^{(1)} U^{(1)})^{-1} dy_5^2 + (U^{(1)} U^{(2)})^{-1} dy_6^2 + dx^\alpha dx^\alpha] \qquad (3.38)$$

4 Depending Functions

Here we consider the ansatzes (2.1) and (2.2) with functions C_a and χ_b which are not independent. We are not going to present a general consideration since it is straightforward but rather involved. We construct only special solutions with depending functions C_a and χ_b. Let us consider the theory (1.1) with $\alpha = 0$, 'electric' + 'magnetic' ansatz with equal number of branches ($E = M$) and $C_a = \chi_a$. Substituting expressions for A, B, and F_i in 'no-force' condition for diagonal part one can obtain

$$-h_a^2 td - v_a^2 ud + v_a^2 \frac{1}{2}\sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{ai} + 1 = 0 \tag{4.39}$$

$$-h_a^2 ud - v_a^2 td + h_a^2 \frac{1}{2}\sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{ai} + 1 = 0 \tag{4.40}$$

These equations are consistent under an assumption

$$v_a^2 = h_a^2 \tag{4.41}$$

Then we have

$$h_a^2 = \frac{2}{d - \sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{ai}} \tag{4.42}$$

A non-diagonal part of the 'no-force' conditions gives

$$\sum_{i=1}^{N} r_i \Delta_{a'i}\Lambda_{ai} - \sum_{i=1}^{N} r_i \Delta_{a'i}\Delta_{ai} - q = 0 \tag{4.43}$$

$$\sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{a'i} - \sum_{i=1}^{N} r_i \Lambda_{a'i}\Lambda_{ai} - s = 0 \tag{4.44}$$

where $a' \neq a$. Note that there is no explicit dependence on D in the conditions (4.43), (4.44).

Some examples. For $D = 11,\ d = 4,\ q = s = 1$ we can put

$$\sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{a'i} = 1,\ \sum_{i=1}^{N} r_i \Delta_{a'i}\Delta_{ai} = 0$$

$$\sum_{i=1}^{N} r_i \Lambda_{a'i}\Lambda_{ai} = 0,\ \sum_{i=1}^{N} r_i \Delta_{ai}\Lambda_{ai} = 0$$

d=3, q=1, s=1

○	○	○						
○			○	○				
○					○	○		
			●		●			●
	●					●		●
		●		●				●

$$ds^2=(H_1H_2H_3)^{\frac{2}{3}}[-(H_1H_2H_3)^{-\frac{4}{3}}dy^2+(H_1^2H_3)^{-2/3}dy_1^2+(H_1^2H_2)^{-2/3}dy_2^2+(H_2^2H_3)^{-2/3}dy_3^2+$$
$$+(H_1H_2^2)^{-2/3}dy_4^2+(H_2H_3^2)^{-2/3}dy_5^2+(H_1H_3^2)^{-2/3}dy_6^2+(H_1H_2H_3)^{-2/3}dy_7^2+dx^\alpha dx^\alpha] \quad (4.45)$$

For the area of horizon we have

$$\mathcal{A}_9=4\pi L^7(H_1H_2H_3)^{2/3} \quad (4.46)$$

Another example. Let us consider $D=10, d=4, q=s=1$

$$\sum_{i=1}^{N} r_i\Delta_{ai}\Lambda_{a'i}=2,\ \sum_{i=1}^{N} r_i\Delta_{a'i}\Delta_{ai}=1$$
$$\sum_{i=1}^{N} r_i\Lambda_{a'i}\Lambda_{ai}=1,\ \sum_{i=1}^{N} r_i\Delta_{ai}\Lambda_{ai}=0$$

$d=4,\ \alpha=0,\ q=1,\ s=1$

○	○	○	○				
○	○			○	○		
○		○		○		○	
○			○		○	○	
				●	●	●	●
		●	●			●	●
	●		●		●		●
	●	●		●			●

We have the following metric

$$ds^2=(H_1H_2H_3H_4)^{\frac{1}{2}}[-(H_1H_2H_3H_4)^{-1}dy^2+(H_1H_2)^{-1}dy_1^2+(H_1H_3)^{-1}dy_2^2+(H_1H_4)^{-1}dy_3^2+$$
$$+(H_2H_3)^{-1}dy_4^2+(H_2H_4)^{-1}dy_5^2+(H_3H_4)^{-1}dy_6^2+dx^\alpha dx^\alpha] \quad (4.47)$$

This solution was discussed in [24].

Acknowledgments

We are grateful to I.V.Volovich for useful discussions and to M.G.Ivanov for attracting our attention to special cases when the stress-energy tensor is non-diagonal. This work is supported by the RFFI grant 96-01-00608.

References

[1] C.Hull and P.Townsend, Nucl.Phys.B438(1995)109

[2] E.Witten, Nucl.Phys.B443(1995)85

[3] P.Townsend, hep-th 9507048

[4] J.H.Schwarz,hep-th/9510086, hep-th/9607201, hep-th/9607201

[5] A.Dubholkar, G.W.Gibbons, J.A.Harvey, F.Ruiz, Nucl.Phys. **340** (1990) 33.

[6] G.T.Horowitz and A.Strominger, Nucl.Phys. B360 (1991) 197.

[7] M.J.Duff and K.S.Stelle, Phys.Lett.**253**(1991) 113.

[8] R.Guven, Phys.Lett.**276**(1992) 49, Phys.Lett.**212**(1988) 277.

[9] R.Kallosh, A.Linde, T.Ortin, A.Peet and A.van Proeyen, Phys.Rev. **D46** (1992) 5278.

[10] H.Lu, C.N.Pope, E.Sergin and K.Stelle, hep-th/9508042.

[11] M.J.Duff, R.R.Khuri and J.X.Lu, Phys.Rep. **259** (1995) 213.

[12] G.W.Gibbons, G.T.Horowitz and P.K.Townsend, Class. Quant. Grav.**12** (1995) 297.

[13] M.Cvetic and D.Youm, hep-th/9510098

[14] H.Lu and C.N.Pope, hep-th/9512153; hep-th/9512012

[15] A.Strominger and C.Vafa, hep-th/9601029

[16] G.T.Horowitz, gr-qc/ 9604051

[17] C.Callan and J.Maldacena, hep-th/9602043

[18] A.A.Tseytlin, hep-th/9601177

[19] G.Papadopoulos and P.K.Townsend, hep-th/9603087

[20] A.A.Tseytlin, hep-th/9604035

[21] H.Lu, C.Pope and K.Stelle, hep-th/9604058

[22] G.Papadopoulos, hep-th/9604068.

[23] G.Lifschytz, hep-th/9604156

[24] I.R.Klebanov and A.A.Tseytlin, hep-th/9604166

[25] K.Behrndt, E.Bergshoeff and B.Janssen, hep-th/9604168

[26] J.P.Gauntlett, D.A.Kastor and J.Traschen, hep-th/9604179.

[27] E.Bergshoeff, R.Kallosh, T.Ortin, hep-th/9605059

[28] A. A. Tseytlin, gr-qc/9608044

[29] A.Volovich, hep-th/9608095

[30] E.Bergshoeff, M.De Roo and S.Panda, hep-th/9609056

[31] G.Papadopoulos and P.K.Townsend, hep-th/9609095

[32] A. A. Tseytlin, hep-th/9609212

[33] I.Ya. Aref'eva, K.S.Viswanathan and I.V.Volovich, hep-th/9609225

[34] K.Behrndt, G;L.Cardoso, B.de Wit, R.Kallosh, D.Lust and T.Mohaupt, hep-th/9610105.

[35] R. Emparan, hep-th/9610170.

[36] C.G.Callan, Jr., S.S.Gubser,I.R.Klebanov and A.A.Tseytlin, hep-th/9610172

[37] I.Ya. Aref'eva and A.Volovich, hep-th/9611026